



Phytochemical and Physicochemical Investigation of *Curcuma Longa* Linn Rhizome

M. A. PAWAR¹, S.S. PATIL^{2,3}, D. M. NAGRIK³

¹Department of chemistry, ShriShivaji Science and Arts College Chikhli, Dist-Buldana

²Department of chemistry, B. B. Arts, N. B. Commerce & B. P. Science College, Digras

³Department of chemistry, G.S. Science and Arts College Khamgaon, Dist-Buldana

Corresponding author: mahesh311286@gmail.com

Abstract

The present article deals with study of phytochemical and physicochemical analysis of *Curcuma longa* Linn. rhizome, a member of family Zingiberaceae. The rhizomes of *Curcuma longa* reported to have good medicinal values in traditional system of medicines. Phytochemical parameters of plant were studied and important chemicals constituents like alkaloids, flavonoids, amino acids, carbohydrate, proteins, saponin and tannins were identified. Physicochemical screening of the rhizome powder showed 11.69% total ash, 0.89% acid insoluble ash, 4.07% water soluble ash, 15.16% water-soluble extractive, 16.10% alcohol soluble extractive and 3.6 pH. This information will be helpful in standardization for quality, purity and sample identification.

Keywords: *Curcuma longa*, physicochemical screening, phytochemical parameters.

Introduction

Plants have been playing an important role in curing the diseases of human beings since time immemorial. The medicinal value of plants is due to some chemically active substances that produce a definite physiological action on the human body. Some important bioactive constituents of plants are alkaloids, tannins and flavonoid and phenolic compounds [1]. These compounds are synthesized by primary or rather secondary metabolism of living organisms. Secondary metabolites are chemically and taxonomically extremely diverse compounds with obscure functions. They are widely used in human therapy, agriculture, scientific research, veterinary and many other areas [2]. Plant products are part of phytomedicines. This can be derived from barks, leaves, flowers, roots, fruits, seeds. Knowledge of the chemical constituents of plants is desirable because such information will be of value for synthesis of complex chemical substances [3, 4 and 5].

According to World Health Organization (WHO), about 80% of individuals from developed countries use traditional medicines, derived from medicinal plants. However, such plants should be investigated to better understand their properties, safety, and efficiency [6]. Plants are used medicinally in different countries and are the source of potential and powerful drugs [7]. Nowadays phytochemical studies have attracted the attention of plant scientists due to the development of new and sophisticated techniques.



Plant synthesizes different types of chemical compounds, which can be differentiated on the basis of their chemical class, functional groups and bio synthetic origin into primary and secondary metabolites [3].

Curcuma longa Linn. belongs to family Zingiberaceae is a perennial herb with pulpy, orange, tuberous roots that grows to about 2 feet in length and is cultivated in India, China, Bangladesh and other Asian countries with a tropical climate. *Curcuma longa* is widely used in Ayurvedic, Unani and Siddha Herbal System. It is recommended for treating diabetes, abdominal pains, menstrual disorder, wounds, eczema, jaundice, inflammations and as a blood purifying activity. Many species of *Curcuma* are traditionally used for their medicinal properties. Antifungal, Antibacterial and Anti inflammatory activity has been reported for species such as *C. long*, *C. zedoria*, *C. aromatica* and *C. amada*[9].

Materials And Methods:

Plant Material Collection

Plant materials of *Curcuma longa* were collected in proper blooming period from Ambabarwa forest area of Buldana district as per the standard method [10]. The plant was identified and authenticated by Dr. Miss V. U. Pochhi, Head, Department of Botany, ShriShivaji Science and Arts College Chikhli. Fresh rhizomes were collected then bring to the laboratory and thoroughly washed with distilled water and shade dried at 28 ± 2 °C. The dried rhizomes were ground well into a fine powder in a mixer grinder. The powder was stored in a polythene bags at room temperatures.

Preparation of the extract

The powder plant material was subjected to hot continuous extraction in a soxhlet apparatus. The powder plant drug was successively extracted with methanol, Acetone, chloroform, Ethyl acetate and hot water. The liquid extracts were collected in tarred conical flask. The solvent was removed by distillation. These extracts were used to study to various qualitative chemical tests and determine the presence of different phytoconstituents like alkaloids, carbohydrates and glycosides, saponins, proteins and amino acid, phenolic compound and tannins, flavonoids etc.

Preliminary Phytochemical Screening

Phytochemical screening of the *C. longa* was done by the standard procedures prescribed by Kokate and Harborne [11, 12].

Physicochemical Evaluation:

The different physicochemical evaluation of the powder rhizome of *c. long* was done to evaluate the quality and purity of the extracted drug. The physicochemical parameters like moisture content, ash values, extractive values, pH etc were determined. [13, 14].



Result and Discussion:

Phytochemical analysis of rhizome of *Curcuma longa* was shown the presence of alkaloids, flavonoids, phenols, steroids, saponins, tannins, glycosides, carbohydrates, proteins, and amino acids while terpenoids found to be absent as given in Table no.1

The similar findings were reported with the extract of rhizome of *C.longaby* earlier workers [15, 16].Phytochemicals evaluation of rhizome in different solvent shows variations in presence of secondary metabolites. Methanolic extract contain 10, chloroform extract 8, acetone extract 8, ethyl acetate extract contain 7 whereas hot water shows 10 numbers of secondary metabolites.

Physicochemical evaluation of *C. longa* rhizome was shown that the results of physicochemical constants found within limit. (Table no. 2) This indicates that the quality and purity of raw material was good enough. The result of moisture content 3.64%w/w implies that the drug is properly dried and stored. The physicochemical parameter like total ash is an important as it shows the purity of drug, which implies presence or absence of foreign material like metallic salts etc. [13]. The physicochemical analysis result for total ash was found to be 11.69w/w. The ash value lies within limit implies purity and quality of crude drug. The water soluble extractive value found to be 15.16%w/w. While ethanol soluble extractive value found to be 16.10%w/w. The water soluble extractive value shows the presence of acids and inorganic compounds[17,14].where as alcohol soluble extractive values represents the presence of polar constituents like phenols, alkaloids, steroids, glycosides, flavonoids [14, 15]. The pH value of crude drug was found to be near about 3.6 which indicate acidic nature of rhizome.

Table 1: Preliminary Phytochemicals Screening of *Curcuma longa* Linn. Rhizome

Sr. No.	Test	Hot water	Methanol	Acetone	Chloroform	Ethyl Acetate
I	Alkaloids					
1	Mayer's Test	+	+	+	-	-
2	Dragandorff's Test	+	+	+	+	+
3	Wagner's Test	+	+	+	-	-
II	Flavonoids					
1	Shinodatest	-	-	-	+	+
2	Lead acetate test	+	+	-	-	-
III	Phenolic compound and Tannins					
1	FeCl ₃ test	+	+	+	-	-



IV	Terpenoids					
1	Liebermann Burchards Test	-	-	-	-	-
V	Steroids					
1	Salkowski test	+	+	+	+	+
VI	Carbohydrates					
	Fehling's Test	+	+	+	+	+
VII	Protein					
	Millon's Test	+	+	-	+	+
VIII	Amino Acid					
	Ninhydrin Test	+	+	+	+	-
IX	Saponins					
	Foam Test	+	+	+	+	-
X	Glucosides					
1	Keller-Killian test	+	+	+	+	+
2	Legal's test	+	+	+	+	+

Where, + = present and - = absent.

Table 2: Physicochemical analysis of *Curcumalonga* Linn. rhizome

Sr. No.	Physicochemical parameters	Values (in %w/w)
1	Total ash value	11.69%
2	Water soluble ash	4.07%
3	Acid insoluble ash	0.89%
4	Moisture content	3.64%
5	Alcohol soluble extractive value	16.10%
6	Water soluble extractive value	15.16%
7	pH	3.6




Fig. 1: Physicochemical parameters of *Curcuma longa* rhizome

Conclusion

The phytochemical screening confirmed the presence of various phytochemical constituents such as alkaloids, flavonoids, amino acids, carbohydrate, proteins, saponin and tannins. Phytochemical constituents confirmed utilization of rhizome for treating diabetes, abdominal pains, menstrual disorder, wounds, eczema, Jaundice, inflammations and as a blood purifying activity. Different Physicochemical parameters such as, total ash, acid insoluble ash, water soluble ash, water soluble extractive and alcohol soluble extractive value were observed. These values can be useful to detect adulteration. All studied standardization parameters like phytochemical screening and physicochemical parameters provide the knowledge in the identification authentication of *Curcuma longa* rhizome.

Reference

- [1] Aiyelaagbe, O. O. and Paul M. Osamudiamen (2009) Phytochemical screening for active compounds in *Mangifera indica* leaves from Ibadan, Oyo State. *Plant science Research*; 2(1):11-13.
- [2] Vasu, K., Goud, J.V., Suryam, A., Singara, Chary, M.A. 2009. Biomolecular and phytochemical analyses of three aquatic angiosperms. *Afr. J. Microbiol. Res.*, 3(8):418-421.
- [3] Mojab, F., M. Kamalinejad, N. Ghaderi and H.R. Vanidipour (2003). Phytochemicals screening of some species of Iranian plants. *Iran. J. Pharm. Res.*, 3: 77-82.
- [4] Parekh, J. and S. Chanda (2007). Antibacterial and phytochemical studies on twelve species of Indian medicinal plants. *Afr. J. Biomed. Res.*, 10: 175-181.
- [5] Parekh, J. and S. Chanda (2008). Phytochemicals screening of some plants from western region of India. *Plant Arch.*, 8: 657-662.



- [6] Arunkumar, S. and Muthuselvam(2009). Analysis of phytochemical constituents and antimicrobial activities of aloe vera L. against clinical pathogens. *World J. Agril. Sc.*, 5(5): 572-576.
- [7] Srivastava, J., J. Lambert and N. Vietmeyer(1996). Medicinal plants: World Bank Technical Paper, 320: 20-27.
- [8] Mojab F., M. Kamalinejad, N. Ghaderi and H. R. Vahidipour(2003). Phytochemical screening of some species of Iranian plants. *Iranian Journal of Pharmaceutical Research*, 2(2):77-82.
- [9] Manimegalai, V, T. Selvaraj and V. Ambikapathy(2011). *Advances in Applied Science Research*, 2. (4):621-628.
- [10] Jain, S.K. and R.R. Rao (1977). *A Handbook of Field and Herbarium Methods*. Today and Tomorrow's Printers and Publishers, New Delhi.
- [11] Kokate, C. K. (1994). *Practical Pharmacognosy*, 4th edn. Vallabh Prakan, New Delhi, 179-181.
- [12] Harbone, J.B and B. L. Turner (1984). *Plant chemosystematics*. Academic press, London. pp: 61-62.
- [13] Kokate, CK. (2007). *Practical Pharmacognosy*. 4th ed., New Delhi; Vallabh Prakashan: pp.1-23.
- [14] Kumar, D., K. Kumar, S. Kumar, T. Kumar, A. Kumar and O. Prakash(2012). *Asian Pacific Journal of Tropical Biomedicine*, 2(3): 169-175.
- [15] Saxena, J. and R. Sahu(2012). Evaluation of Phytochemical constituents in Conventional and Non-conventional species of Curcuma. *International Research Journal of Pharmacy*, 3(8):203-204.
- [16] Swadhini, S.P., R. Santosh, C. Uma, S. Mythili and A. Sathivelu(2011). Phytochemical Screening and Antimicrobial activity of five medicinal plants against *Myrothecium SP*. *International Journal of Pharma and Biosciences*, 2(1): B272-B279.
- [17] Anonymous. (1996). *Indian Pharmacopoeia, Vol-II*, Ministry of Health and Family welfare, Govt of India, New Delhi; Controller of Publications, A- 53-54, A-95, A-97, A-109